



Synonym

CCR5,CMKBR5,CD195,CC-CKR-5,CHEMR13

Source


Human CCR5 Full Length Protein-VLP(CC5-H52P3) is expressed from human 293 cells (HEK293). It contains AA Met 1 - Leu 352 (Accession # [P51681-1](#)).

Predicted N-terminus: Met 1

Molecular Characterization

The protein has a calculated MW of 40.5 kDa | 56.1 kDa.

Virus-like particles(VLPs) are formed by self-assembly of envelop/capsid proteins from viruses. Membrane Proteins can be constituted in-situ with VLPs produced from HEK293 cell cultures. These VLPs concentrate conformationally intact membrane proteins directly on the cell surface and produce soluble, high-concentration proteins perfect for immunization and antibody screening.



The VLPs provide the display of properly folded membrane proteins in their native cellular membrane in a compact size of 100~300 nm diameter (similar to the size of most viruses) making it optimal targets for dendritic cells in vivo and surface attachment for phage display.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

*The isotype control of empty/mock VLP (Cat. No. [VLP-N5213](#)) is sold separately and not included in protein, you can follow [this link](#) for product information.

Formulation

The VLPs are highly immunogenic, so the immunization strategy should be optimized (antigen dose, regimen and adjuvant).

Supplied as 0.2 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Shipping

This product is supplied and shipped with dry ice, please inquire the shipping cost.


Storage

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- The product MUST be stored at -70°C or lower upon receipt;
- -70°C for 12 months under sterile conditions.

Bioactivity-ELISA




Immobilized Human CCR5 Full Length Protein-VLP (Cat. No. CC5-H52P3) at 2 µg/mL (100 µL/well) can bind Anti-Human CCR5 Antibody, Mouse IgG2a with a linear range of 2-78 ng/mL (QC tested).

Discounts, Gifts,
and more!





Identity-DLS



The mean peak Radius of VLP is 55-75 nm with more than 95% intensity as determined by dynamic light scattering (DLS).

Background

Receptor for a number of inflammatory CC-chemokines including CCL3/MIP-1-alpha, CCL4/MIP-1-beta and RANTES and subsequently transduces a signal by increasing the intracellular calcium ion level. May play a role in the control of granulocytic lineage proliferation or differentiation.⁵ Publications (Microbial infection) Acts as a coreceptor (CD4 being the primary receptor) of human immunodeficiency virus-1/HIV-1.

Clinical and Translational Updates

Discounts, Gifts,
and more!

