



**Synonym**

IL-23 alpha & IL-12 beta

**Source**

Human IL-23A&Rat IL-12B Heterodimer Protein, His Tag&Tag Free(ILB-HR52W3) is expressed from human 293 cells (HEK293). It contains AA Arg 20 - Pro 189 (IL23A) & Met 23 - Ser 335 (IL12B) (Accession # [Q9NPF7-1](#) (IL23A) & [Q9R278-1](#) (IL12B)).

Predicted N-terminus: Arg 20 (IL23A) & Met 23 (IL12B)

**Molecular Characterization**

Poly-his	IL23A (Arg 20 - Pro 189) Q9NPF7-1
	IL12B (Met 23 - Ser 335) Q9R278-1

Human IL-23A & Rat IL-12B Heterodimer Protein, His Tag&Tag Free, produced by co-expression of IL-23A and IL-12B, has a calculated MW of 20.6 kDa (IL-23A) and 35.9 kDa (IL-12B). Subunit IL-23A is fused with a polyhistidine tag at the N-terminus and subunit IL-12B contains no tag. The reducing (R) protein migrates as 23 kDa (IL-23A) and 45 kDa and 47 kDa (IL-12B) respectively due to glycosylation.

**Endotoxin**

Less than 1.0 EU per µg by the LAL method.

**Purity**

>90% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

**Formulation**

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

**Reconstitution**

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

**Storage**


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


**SDS-PAGE**



Human IL-23A&Rat IL-12B Heterodimer Protein, His Tag&Tag Free on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

**Bioactivity-ELISA**

**SEC-MALS**



The purity of Human IL-23A&Rat IL-12B Heterodimer Protein, His Tag&Tag Free (Cat. No. ILB-HR52W3) is more than 90% and the molecular weight of this protein is around 65-75 kDa verified by SEC-MALS.

[Report](#)


Discounts, Gifts,  
and more!






Human IL-23A & Rat IL-12B Heterodimer Protein, His Tag&Tag Free ELISA

0.5 µg of Human IL-23 R, Fc Tag per well



Human IL-23A & Rat IL-12B Heterodimer Protein, His Tag&Tag Free ELISA

0.5 µg of Human IL-23A & Rat IL-12B Heterodimer Protein, His Tag&Tag Free per well



Immobilized Human IL-23 R, Fc Tag (Cat. No. ILR-H5254) at 5 µg/mL (100 µL/well) can bind Human IL-23A & Rat IL-12B Heterodimer Protein, His Tag&Tag Free (Cat. No. ILB-HR52W3) with a linear range of 0.039-0.156 µg/mL (QC tested).

Immobilized Human IL-23A & Rat IL-12B Heterodimer Protein, His Tag&Tag Free (Cat. No. ILB-HR52W3) at 5 µg/mL (100 µL/well) can bind Human IL-12 R beta 1, Fc Tag (Cat. No. ILB-H5255) with a linear range of 5-78 ng/mL (Routinely tested).

## Background

Interleukin-23 subunit alpha (IL-23 alpha) can associate with IL12B to form the IL-23 interleukin, a heterodimeric cytokine which functions in innate and adaptive immunity. IL-23 may constitute with IL-17 an acute response to infection in peripheral tissues. IL-23 binds to a heterodimeric receptor complex composed of IL12RB1 and IL23R, activates the Jak-Stat signaling cascade, stimulates memory rather than naive T-cells and promotes production of proinflammatory cytokines. IL-23 induces autoimmune inflammation and thus may be responsible for autoimmune inflammatory diseases and may be important for tumorigenesis.

## Clinical and Translational Updates

Discounts, Gifts,  
and more!

