



Synonym

LYPD1, Ly6/PLAUR domain-containing protein 1, FLJ41033, LY6/PLAUR domain containing 1, LYPDC1, PHTS, Putative HeLa tumor suppressor

Source

Human LYPD1, Fc Tag (LY1-H5254) is expressed from human 293 cells (HEK293). It contains AA Leu 21 - Gly 115 (Accession # [Q8N2G4-1](#)). Predicted N-terminus: Leu 21

Molecular Characterization

LYPD1(Leu 21 - Gly 115) Q8N2G4-1	Fc(Pro 100 - Lys 330) P01857
-------------------------------------	---------------------------------

This protein carries a human IgG1 Fc tag at the C-terminus.

The protein has a calculated MW of 36.8 kDa. The protein migrates as 43 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

SDS-PAGE




Human LYPD1, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With [Star Ribbon Pre-stained Protein Marker](#)).

Bioactivity-ELISA

LYPD1 ELISA

Immobilized Human LYPD1, Fc Tag (Cat. No. LY1-H5254) at 1 µg/mL (100 µL/well) can bind Anti-LYPD1 antibody with a linear range of 0.02-0.625 µg/mL (QC tested).

SEC-MALS



The purity of Human LYPD1, Fc Tag (Cat. No. LY1-H5254) is more than 85% and the molecular weight of this protein is around 142-172 kDa verified by SEC-MALS.

[Report](#)

Discounts, Gifts,
and more!





Background

Believed to act as a modulator of nicotinic acetylcholine receptors (nAChRs) activity. In vitro increases receptor desensitization and decreases affinity for ACh of alpha-4:beta-2-containing nAChRs. May play a role in the intracellular trafficking of alpha-4:beta-2 and alpha-7-containing nAChRs and may inhibit their expression at the cell surface. May be involved in the control of anxiety.

Clinical and Translational Updates

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.

Discounts, Gifts,
and more!

