



### Synonym

TNFRSF4,OX40,CD134,OX40L receptor,ACT35,TXGP1L

### Source

Marmoset OX40, His Tag(OX0-M522b) is expressed from human 293 cells (HEK293). It contains AA Leu 29 - Ala 214 (Accession # [F7DHA3-1](#)).

Predicted N-terminus: Leu 29

### Molecular Characterization

OX40(Leu 29 - Ala 214)  
F7DHA3-1 Poly-his

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 21.8 kDa. The protein migrates as 42-45 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

### Endotoxin

Less than 1.0 EU per µg by the LAL method.

### Purity

>90% as determined by SDS-PAGE.

### Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

### Reconstitution

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

### Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

### SDS-PAGE





Marmoset OX40, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 90%.

### Bioactivity-ELISA


Discounts, Gifts,  
and more!





Immobilized Marmoset OX40, His Tag (Cat. No. OX0-M522b) at 2 µg/mL (100 µL/well) can bind Human OX40 Ligand, Fc Tag, premium grade (Cat. No. OXL-H5266) with a linear range of 0.2-31 (QC tested).

Immobilized Marmoset OX40, His Tag (Cat. No. OX0-M522b) at 2 µg/mL (100 µL/well) can bind Rhesus macaque OX40 Ligand, Mouse IgG2a Fc Tag (Cat. No. OXL-R5259) with a linear range of 0.2-4 ng/mL (Routinely tested).



Immobilized Marmoset OX40, His Tag (Cat. No. OX0-M522b) at 1 µg/mL (100 µL/well) can bind Anti-OX40 Antibody, Human IgG2 with a linear range of 4-63 ng/mL (Routinely tested).

**Background**

Tumor necrosis factor receptor superfamily member 4 (TNFRSF4) is also known as ACT35 antigen, OX40L receptor, TAX transcriptionally-activated glycoprotein 1 receptor, CD antigen CD134, OX40. OX40 / TNFRSF4 contains four TNFR-Cys repeats. TNFRSF4 is receptor for TNFSF4 / OX40L / GP34 and can interact with TRAF2, TRAF3 and TRAF5.

**Clinical and Translational Updates**

