

Synonym

Spike,S protein RBD,Spike glycoprotein Receptor-binding domain,S glycoprotein RBD,Spike protein RBD

Source

SARS-CoV-2 S protein RBD, Fc Tag(SPD-C5255) is expressed from human 293 cells (HEK293). It contains AA Arg 319 - Phe 541 (Accession # [QHD43416.1](#)). Predicted N-terminus: Arg 319

Molecular Characterization

S protein RBD(Arg 319 - Phe 541) QHD43416.1	Fc(Pro 100 - Lys 330) P01857
--	---------------------------------

This protein carries a human IgG1 Fc tag at the C-terminus

The protein has a calculated MW of 51.5 kDa. The protein migrates as 60-65 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



SARS-CoV-2 S protein RBD, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.



Bioactivity-ELISA

SEC-MALS



The purity of SARS-CoV-2 S protein RBD, Fc Tag (Cat. No. SPD-C5255) is more than 90% and the molecular weight of this protein is around 110-125 kDa verified by SEC-MALS.


[Report](#)




Immobilized Human ACE2, His Tag (Cat. No. AC2-H52H8) at 2 µg/mL (100 µL/well) can bind SARS-CoV-2 S protein RBD, Fc Tag (Cat. No. SPD-C5255) with a linear range of 0.2-5 ng/mL (QC tested).

Immobilized SARS-CoV-2 S protein RBD, Fc Tag (Cat. No. SPD-C5255) at 2 µg/mL (100 µL/well) can bind Anti-SARS-CoV-2 Neutralizing Antibody, Human IgG1 (Cat. No. SAD-S35) with a linear range of 0.4-3 ng/mL (Routinely tested).

Bioactivity-BLI



Loaded SARS-CoV-2 S protein RBD, Fc Tag (Cat. No. SPD-C5255) on Protein A Biosensor, can bind Human ACE2, His Tag (Cat. No. AC2-H52H8) with an affinity constant of 2.16 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).



Loaded SARS-CoV-2 S protein RBD, Fc Tag (Cat. No. SPD-C5255) on Protein A Biosensor, can bind Cynomolgus ACE2, His Tag (Cat. No. AC2-C52H7) with an affinity constant of 4.48 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).

Background

It's been reported that Coronavirus can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

Clinical and Translational Updates

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.