



Synonym

TNFRSF1B,CD120b,TBPII,TNF-R-II,TNF-R75,TNFBR,TNFR1B,TNFR2,TNFR80,p75,p75TNFR

Source

Human TNFR2, His Tag(TN2-H5227) is expressed from human 293 cells (HEK293). It contains AA Leu 23 - Asp 257 (Accession # [AAA36755](#)).

Predicted N-terminus: Leu 23

Molecular Characterization

TNFR2(Leu 23 - Asp 257)
AAA36755 Poly-his

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 26.0 kDa. The protein migrates as 35-45 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Human TNFR2, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With [Star Ribbon Pre-stained Protein Marker](#)).

Bioactivity-ELISA

SEC-MALS




The purity of Human TNFR2, His Tag (Cat. No. TN2-H5227) is more than 85% and the molecular weight of this protein is around 35-47 kDa verified by SEC-MALS.


[Report](#)

Discounts, Gifts,
and more!





Immobilized Human TNF-alpha, premium grade (Cat. No. TNA-H4211) at 5 µg/mL (100 µL/well) can bind Human TNFR2, His Tag (Cat. No. TN2-H5227) with a linear range of 5-39 ng/mL (QC tested).



Immobilized ActiveMax® Human TNF-beta, Tag Free at 5 µg/mL (100 µL/well) can bind Human TNFR2, His Tag (Cat. No. TN2-H5227) with a linear range of 2-62 ng/mL (Routinely tested).

Background

TNF RI is also known as the p60 or p55 TNFR) and TNF RII (the p75 or p80 TNFR) are two distinct type I transmembrane glycoproteins that bind TNF with high affinity. Both RI and RII are prototypic members of the TNF receptor superfamily and have been designated TNFRSF1A and TNFRSF1B, respectively. Human TNF RII cDNA encodes a 461 amino acid (aa) residue precursor protein with a 22 aa putative signal peptide, a 235 aa extracellular domain, a 20 aa transmembrane domain and a 174 aa cytoplasmic domain. TNFRII is expressed in fetal brain. The protein is produced naturally as a soluble form (sTNFRII). The soluble receptor inhibits TNF α action by competing with cell surface receptors in binding TNF α , thereby blocking its biologic effects. TNFRII is strongly expressed at the cartilage-pannus junction, and plays a major role in a subset of families with multiple cases of rheumatoid arthritis (RA). Further, high plasma levels of sTNFRII were significantly associated with increased incidence of coronary heart disease, independent of established cardiovascular risk factors, and seems to be useful for monitoring the inflammatory activity of sarcoidosis.

Clinical and Translational Updates

Discounts, Gifts,
and more!

